



SOME DATA ON SNOW AND THEIR CLIMATE ANALYSIS, IN THE WESTERN AREA OF LAKE MICHIGAN: CASE STUDY

Hysen MANKOLLI,

¹Expert in Ecology, Plainfield, Illinois, USA; Consultant Scientific Earth System Science Interdisciplinary Center (ESSIC), University of Maryland College Park, USA; Chief Editor of IJEES journal;
Email: hysenmankolli@yahoo.com;

Abstract

Some data on snow and their climate analysis, in the western area of lake Michigan: Case study. The data obtained refer to the stations Chicago, Freeport, Rockford, Kenosha, Aurora, Gary, Joliet, Ottawa, Dixon, Clinton, Dekalb, Kankakee, Pontiac, Peoria, Bloomington.

The climate of the Chicago Area is with cold winters and scattered snowfall in the winter months. The study shows some estimates of snowfall trends over a multi-year period. Data refers to reports from the National Oceanic and Atmospheric Administration National Weather Service, Chicago, IL, 333 West University Drive, Romeoville, IL. The analysis of snow precipitation amount data is based on the application of the mathematical method Excel.

The results show that precipitation values are different over the years. Trends are given by regression equations. For January 1985-2021: $y = 4.4576x + 1937.5$, $R^2 = 0.0378$. The tendency of the value of snowfall for the months of January is with a constant 4.4576. In the period Seasonal Snowfall Totals for Chicago from 1884 to 2021: $y = 2.3353x + 1890.2$, $R^2 = 0.0868$. The tendency of the value of snowfall for each year period is with a constant 2.3353. The smallest amount of snow precipitation is estimated for the years 1920-1921 with 9.8 inches. The highest amount of snow precipitation is estimated for the years 1978-1979 with 89.7 inch. All result are based on statistical method. Our statewide snowfall records in Illinois start in 1902. There is no long-term trend in snowfall since 1902.

However, some decades were snowier than others, such as the 1910s, 1960s, and 1970s. In fact, the 1970s were the snowiest decade on record with an average snowfall of 27.2 inches. Snowfall amounts dropped steeply with less year to year variability for much of the 1980s and into the early 2000s. However, snowfall amounts in the last 6 winters have been more variable with the winter of 2014 being about as snowy as the late 1970s. The snowiest winter on record was 1979 with 44.5 inches. Here are the 5 snowiest winters on record: 1979 with 44.5 inches; 1978 with 44.4 inches; 1912 with 39.5 inches; 2014 with 39.4 inches; 1960 with 38.6 inches;

Snowfall is accumulated from July 1 of the first year to June 30 of the second year. The second year is used in the plot and table (for example, 2006 refers to the 2005-2006 season, source: state climatologist Illinois).

Introduction

Snow is an important element of the climate. Many studies on snowfall have been conducted in the USA and Europe. Regardless of the results achieved, studies and trends on snow performance, time and quantity, always remain to be evaluated.

In each case of evaluation are considered the time of snowfall, the amount of snowfall, monthly, annual multi-year period as well as mathematical and statistical methods.

Material and methods

The analysis of snow precipitation amount data is based on the application of the mathematical method Excel. The results show that precipitation values are different over the years. Trends are given by regression equations.

Data

The data obtained refer to the stations Chicago, Freeport, Rockford, Kenosha, Aurora, Gary, Joliet, Ottawa, Dixon, Clinton, Dekalb, Kankakee, Pontiac, Peoria, Bloomington. The climate of the Chicago Area is with cold winters and scattered snowfall in the winter months.

The study shows some estimates of snowfall trends over a multi-year period. Data refers to reports from the National Oceanic and Atmospheric Administration National Weather Service, Chicago, IL, 333 West University Drive, Romeoville, IL,
https://www.weather.gov/lot/Chicago_seasonal_snow.

RESULTS

After analyzing datas, for snowfall November data in Chicago area based at figure 1, results:

- Snow inch for each year at every November it is different, started with 0 inch 1968, 1 inch 1884, 1896, 1902, 1908, 1938, 1944, 1950, 1956, and 3 and 4 inches in 1896, 1968 and maximum inches in 2001 with 12.

After analyzing datas, for seasonal snow in Chicago area based at figure 2, results:

- The tendency of the regression equation is positive.
- The coefficient for each year is 2,353.
- The functional bond is weak with $R^2 = 0.08$.

After analyzing datas, for smallest and largest snowfall data in Chicago area based at figure 3, results:

- The tendency of the regression equation is almost linear.
- The coefficient for each year is 4.7683.
- The functional bond is weak with $R^2 = 0.009$.




Figure 3. Chicago Monthly Snowfall Amounts, January, Normal 11.3 Inches

CONCLUSIONS

The analysis of snow precipitation amount data is based on the application of the mathematical method Excel. The results show that precipitation values are different over the years. Trends are given by regression equations. For January 1985-2021: $y = 4.4576x + 1937.5$, $R^2 = 0.0378$. The tendency of the value of snowfall for the months of January is with a constant 4.4576. In the period Seasonal Snowfall Totals for Chicago from 1884 to 2021: $y = 2.3353x + 1890.2$, $R^2 = 0.0868$. The tendency of the value of snowfall for each year period is with a constant 2.3353. The smallest amount of snow precipitation is estimated for the years 1920-1921 with 9.8 inches. The highest amount of snow precipitation is estimated for the years 1978-1979 with 89.7 inch. All result are based on statistical method.

Acknowledgement. This paper has been prepared a part of independent Study CLIMATIC DATA AND SNOW DEPTH IN SOME REGIONS AT USA AND ALBANIA, project 2023 no. 02012023.

Conflicts of interest. The authors declare no conflicts of interest.

REFERENCES

- Cezar Kongoli¹², 2016. LARGE - SCALE SNOW DEPTH – ELEVATION RELATIONSHIPS OF WORLD'S MAJOR MOUNTAIN REGIONS, International Journal of Ecosystems and Ecology Science (IJEES): Vol. 6 (4): 653-658 (2016);
- Cezar Kongoli¹², Robert J. Kuligowski², Sean Helfrich², Peter Romanov³², 2014. MONITORING PRECIPITATION AND SNOW COVER USING MULTI-SENSOR SATELLITE AND IN-SITU DATA, International Journal of Ecosystems and Ecology Science (IJEES); Vol. 4 (2): 293-300 (2014);
- Cezar Kongoli¹², Sean Helfrich², Robert J. Kuligowski², 2015. SATELLITE-BASED ESTIMATION OF HYDROLOGIC COMPONENTS – APPLICATION TO SNOW AND PRECIPITATION, International Journal of Ecosystems and Ecology Science (IJEES): Vol. 5 (2): 261- 268 (2015);
- Cezar Kongoli¹², Yu Zhang³, Zhengtao Cui⁴², Robert J. Kuligowski², 2017. ESTIMATION OF SNOW AND RAINFALL FOR HYDROLOGICAL ANALYSIS OF A MID-WINTER FLOODING EVENT IN WESTERN UNITED STATES, International Journal of Ecosystems and Ecology Science (IJEES): Vol. 7 (4): 651-656 (2017);
- Hall, D.K., G.A. Riggs, V.V. Salomonson, N.E. DiGirolamo and K.A. Bayr, 2002. MODIS snow-cover products, Remote Sensing of Environment, 83:181-194;
- Hysen Mankolli¹², Andi Mankolli^{3*}, 2017. SNOW AND RAIN PRECIPITATION IN SOME REGIONS OF ALBANIA, International Journal of Ecosystems and Ecology Science (IJEES): Vol. 7 (3): 645-650 (2017);
- Kongoli, C. E. and Bland, W. L.(2000) : Long-term snow depth simulations using a modified atmosphere-land exchange model, Agr. Forest Meteorol., 104, 273-287;